

Khovanov homology of graph-links

Igor Nikonov

Abstract

We define Khovanov homology over $\mathbb{Z}_2$ for graph-links.

1 Introduction

Graph-links [4, 5] are combinatorial analog of classical and virtual links. "Diagrams" of graph-links are simple undirected labeled graphs and graph-links themselves are equivalence classes of the graphs modulo formal Reidemeister moves. In knot theory graph-links appear as intersection graphs of rotating chord diagrams of links. It is known that intersection graph determines chord diagram uniquely up to mutations [2]. Thus any link invariant that does not distinguish mutant links is a candidate for an invariant of graph-links. For example, Alexander polynomial, Jones polynomial and HOMFLY are of such type. The problem is the equivalence relation of the graph-links can be too strong so there can exist two chord diagrams which have different values of the invariant but whose intersection graphs coincide as graph-links. So the problem of finding a natural construction that would extend the invariant from intersection graphs to graph-links is nontrivial. For Jones polynomial this problem was solved by D. Ilyutko and V. Manturov [4]. This article provides a solution for (odd) Khovanov homology with coefficients in $\mathbb{Z}_2$.

A theory analogous to the theory of graph-links was developed by L. Traldi and L. Zulli [7]. Their approach (called looped interlacement graphs theory) is based on Gauss diagrams of knots instead of rotating chord diagrams. D. Ilyutko showed these two theories are closely related [3]. So the construction of the Khovanov homology described below will provide as well (after the necessary reformulation) a definition of the Khovanov homology for the looped interlacement graphs.

Odd Khovanov homology of links was defined by P. Ozsváth, J. Rasmussen and Z. Szabó [6]. J. Bloom proved that odd Khovanov homology is mutant invariant [1] and we can expect it to be extended to an invariant of graph-links. In fact the construction of J. Bloom over $\mathbb{Z}_2$ can be directly transferred to the graph-link theory. One can also consider this invariant as the ordinary Khovanov homology of graph-links because for links both homology theories coincide modulo 2. On the other hand, integer-valued Khovanov homology is not mutant invariant [8] so we don't anticipate any adaption of it for graph-links.

2 Graph-links

Let G be an undirected graph without loops and multiple edges and $\mathcal{V} = \mathcal{V}(G)$ be the set of its vertices. We assume G be a *labeled* graph, i.e. every vertex v in G is endowed with a pair (a, α) where $a \in \{0, 1\}$ is the *framing* $\text{fr}(v)$ of v and $\alpha \in \{-1, 1\}$ is the *sign* $\text{sgn}(v)$ of the vertex v .

Let us fix an enumeration of vertices of G . We define the *adjacency matrix* $A(G) = (a_{ij})_{i,j=1,\dots,n}$ over $\mathbb{Z}_2$ as follows: $a_{ij} = 1$ if and only if the vertices v_i and v_j are adjacent, and $a_{ij} = 0$ otherwise. Besides we set $a_{ii} = \text{fr}(v_i)$.

Let $v \in \mathcal{V}$. The set of all vertices in $\mathcal{V}$ adjacent to v is called the *neighbourhood* of the vertex v and denoted $N(v)$.

Let us define the Reidemeister moves of labeled graphs.

Ω_1 . The first Reidemeister move is an addition/removal of an isolated vertex labeled $(0, \pm 1)$.

Ω_2 . The second Reidemeister move is an addition/removal of two adjacent (resp. nonadjacent) vertices having the labels $(1, -1)$ and $(1, 1)$ (resp. $(0, -1)$ and $(0, 1)$) and the same neighbourhoods.

Ω_3 . The third Reidemeister move is defined as follows. Let u, v, w be three vertices of G having the labels $(0, -1)$ and u be adjacent only to v and w . Then we only change the adjacency of u with the vertices $t \in N(v) \setminus N(w) \cup N(w) \setminus N(v)$ (for other pairs of vertices we don't change their adjacency). In addition, we change the signs of v and w to '+1'. The inverse operation is also called the third Reidemeister move.

Ω_4 . The fourth Reidemeister move is defined as follows. We take two adjacent vertices u labeled $(0, \alpha)$ and v labeled $(0, \beta)$. Then we change the label of u to $(0, -\beta)$ and the label of v to $(0, -\alpha)$. Besides, we change the adjacency for each pair (t, w) of vertices where $t \in N(u)$ and $w \in N(v) \setminus N(u)$ or $t \in N(v)$ and $w \in N(u) \setminus N(v)$.

Ω'_4 . In this fourth move we take a vertex v with the label $(1, \alpha)$. We change the adjacency for each pair (t, w) of vertices where $t, w \in N(u)$. Besides, we change the sign of v and the framing for each $w \in N(u)$.

Definition 1. A *graph-link* is the equivalence class of a simple labeled graph modulo $\Omega_1 - \Omega'_4$ moves.

3 Khovanov homology of graph links

Let G be a simple labeled graph with n vertices and $A = A(G)$ be its adjacency matrix. We now formulate mod 2 version of Bloom's construction of odd Khovanov homology. The reason why we use $\mathbb{Z}_2$ is the chain complex construction depends on the coefficients of the adjacency matrix A and the matrix A is defined over $\mathbb{Z}_2$.

Suppose $s \subset \mathcal{V} = \mathcal{V}(G)$. We shall call subsets of $\mathcal{V}$ as *states*. Define $G(s)$ to be the complete subgraph in G with the set of vertices s and denote $A(s) =$

$A(G(s))$. Consider the vector space

$$V(s) = \mathbb{Z}_2 \langle x_1, \dots, x_n \mid r_1^s, \dots, r_n^s \rangle$$

where relations $r_1^s, \dots, r_n^s$ are given by the formula

$$r_i^s = \begin{cases} x_i + \sum_{\{j \mid v_j \in s\}} a_{ij} x_j, & \text{if } v_i \notin s, \\ \sum_{\{j \mid v_j \in s\}} a_{ij} x_j, & \text{if } v_i \in s \end{cases} \quad (1)$$

The dimension of $V(s)$ is equal to $\text{corank } A(s)$.

There is a natural bijection between states $s \subset \mathcal{V}$ and vertices of the hypercube $\{0, 1\}^n$: the state s corresponds to the vector $(\alpha_1, \dots, \alpha_n)$ where $\alpha_i = 0$ if $v_i \in s$ and $\text{sgn}(v_i) = 1$ or if $v_i \notin s$ and $\text{sgn}(v_i) = -1$ and $\alpha = 1$ otherwise. Every edge of the hypercube is of the type $s \rightarrow s \oplus i$ where $s \oplus i$ denotes $s \cup \{v_i\}$ if $v_i \notin s$ and $s \setminus \{v_i\}$ if $v_i \in s$. We orient the arrow so that $v_i \notin s$ if $\text{sgn}(v_i) = -1$ and $v_i \in s$ if $\text{sgn}(v_i) = 1$.

We assign to every edge $s \rightarrow s \oplus i$ the map $\partial_{s \oplus i}^s : \bigwedge^* V(s) \rightarrow \bigwedge^* V(s \oplus i)$ of exterior algebras defined by the formula

$$\partial_{s \oplus i}^s(u) = \begin{cases} x_i \wedge u & \text{if } x_i = 0 \in V(s), \\ u & \text{if } x_i \neq 0 \in V(s). \end{cases} \quad (2)$$

Consider the chain complex

$$C(G) = \bigoplus_{s \subset \mathcal{V}} \bigwedge^* V(s)$$

with differential

$$\partial(u) = \sum_{\{s, s' \subset \mathcal{V} \mid s \rightarrow s'\}} \partial_{s'}^s(u).$$

Proposition 1. *Chain complex $(C(G), \partial)$ is well defined.*

Proof. We need to show that the maps $\partial_{s \oplus i}^s$ are well defined and that each 2-face in the hypercube of states is commutative.

Lemma 1. *Let us consider a state s and an index i such that $v_i \notin s$. We can assume without loss of generality that $A(s \oplus i) = \begin{pmatrix} A & a^\top \\ a & \alpha \end{pmatrix}$ where $A = A(s)$. Then*

1. $x_i = 0 \in V(s)$ iff $\text{rank } A = \text{rank} \begin{pmatrix} A \\ a \end{pmatrix}$;
2. $x_i = 0 \in V(s \oplus i)$ iff $\text{rank} \begin{pmatrix} A \\ a \end{pmatrix} + 1 = \text{rank} \begin{pmatrix} A & a^\top \\ a & \alpha \end{pmatrix}$.

Proof. It follows from the relation $x_i = \sum_{\{j \mid v_j \in s\}} a_{ij}x_j$ that the equality $x_i = 0 \in$

$V(s)$ means the vector a depends on the rows of the adjacency matrix A . This leads to the first statement of the lemma.

The equality $x_i = 0 \in V(s \oplus i)$ implies that the vector $(0 \ 1)$ depends on the rows of the matrix $A(s \oplus i)$. Then $\text{rank} \begin{pmatrix} A & a^\top \\ a & \alpha \end{pmatrix} = \text{rank} \begin{pmatrix} A & a^\top \\ a & \alpha \\ 0 & 1 \end{pmatrix}$. But

$$\text{rank} \begin{pmatrix} A & a^\top \\ a & \alpha \\ 0 & 1 \end{pmatrix} = \text{rank} \begin{pmatrix} A & 0 \\ a & 0 \\ 0 & 1 \end{pmatrix} = \text{rank} \begin{pmatrix} A \\ a \end{pmatrix} + 1.$$

□

Lemma 2. For any symmetric matrix A if $\text{rank} \begin{pmatrix} A & a^\top \\ a & \alpha \end{pmatrix} = \text{rank } A + 1$ then $\text{rank} \begin{pmatrix} A \\ a \end{pmatrix} = \text{rank } A$.

Proof. Assume that $\text{rank} \begin{pmatrix} A \\ a \end{pmatrix} \neq \text{rank } A$. Then $\text{rank} \begin{pmatrix} A & a^\top \\ a & \alpha \end{pmatrix} = \text{rank} \begin{pmatrix} A \\ a \end{pmatrix}$ and the vector $\begin{pmatrix} a^\top \\ \alpha \end{pmatrix}$ depends on the columns of the matrix $\begin{pmatrix} A \\ a \end{pmatrix}$. Then the vector $a^\top$ depends of the columns of A and (after transposition) the vector a depends on the rows of A . Hence, $\text{rank } A = \text{rank} \begin{pmatrix} A \\ a \end{pmatrix}$ that contradicts to the initial assumption. □

Corollary 1. For any state s and any index i we have

1. $\dim V(s \oplus i) = \dim V(s) + 1$ iff $x_i = 0 \in V(s)$ and $x_i \neq 0 \in V(s \oplus i)$;
2. $\dim V(s \oplus i) = \dim V(s) - 1$ iff $x_i \neq 0 \in V(s)$ and $x_i = 0 \in V(s \oplus i)$;
3. $\dim V(s \oplus i) = \dim V(s)$ iff $x_i = 0 \in V(s)$ and $x_i = 0 \in V(s \oplus i)$.

The case $x_i \neq 0 \in V(s)$ and $x_i \neq 0 \in V(s \oplus i)$ is impossible.

Proof. The statements of the corollary follow from Lemmas 1,2 and the fact $\dim V(s) = \text{corank } A(s)$, $\dim V(s \oplus i) = \text{corank } A(s \oplus i)$. □

We call the first two cases in the proposition *even* and the third case *odd*. From the definition of the differentials we have $\partial_{s \oplus i}^s = 0$ in the odd case.

Correctness of chain maps.

Let us consider the map $\partial_{s \oplus i}^s : \bigwedge^* V(s) \rightarrow \bigwedge^* V(s \oplus i)$. We must check that the relations are mapped to relations, that is for any element u and any index j there exist elements $u_k \in V(s \oplus i)$ such that

$$\partial_{s \oplus i}^s(r_j^s \wedge u) = \sum_k r_k^{s \oplus i} \wedge u_k \in V(s \oplus i).$$

For any j we have $r_j^s = r_j^{s \oplus i} + \alpha_j x_i$ for some $\alpha_j \in \mathbb{Z}_2$. If $x_i = 0 \in V(s \oplus i)$ then

$$\partial_{s \oplus i}^s(r_j^s \wedge u) = r_j^s \wedge u' = r_j^{s \oplus i} \wedge u' + \alpha_j x_i \wedge u' = r_j^{s \oplus i} \wedge u'$$

in $V(s \oplus i)$, where $u' = u$ or $u' = 0$. If $x_i \neq 0 \in V(s \oplus i)$ then $x_i = 0 \in V(s)$ and

$$\partial_{s \oplus i}^s(r_j^s \wedge u) = x_i \wedge r_j^s \wedge u = x_i \wedge r_j^{s \oplus i} \wedge u + \alpha_j x_i \wedge x_i \wedge u = r_j^{s \oplus i} \wedge (x_i \wedge u).$$


In any case the map $\partial_{s \oplus i}^s$ is well defined.

Commutativity of 2-faces.

Every 2-face of hypercube looks like

$$\begin{array}{ccc} \bigwedge^* V(s \oplus j) & \xrightarrow{\partial_{s \oplus i \oplus j}^{s \oplus j}} & \bigwedge^* V(s \oplus i \oplus j) \\ \partial_{s \oplus j}^s \uparrow & & \uparrow \partial_{s \oplus i \oplus j}^{s \oplus i} \\ \bigwedge^* V(s) & \xrightarrow{\partial_{s \oplus i}^s} & \bigwedge^* V(s \oplus i). \end{array}$$

According to dimensions of spaces $V(s')$, $s' = s, s \oplus i, s \oplus j, s \oplus i \oplus j$ we have five types of diagrams without odd edges:



Here the number at the place of the state s' is equal to $\dim V(s') - \dim V(s) = \text{corank } A(s') - \text{corank } A(s)$ and the label $z = 1, x_i, x_j$ at the arrow for the map $\partial_{s'}^{s''}$ means that $\partial_{s'}^{s''}(u) = z \wedge u$.

2-faces of types 1,2,3 are obviously commutative. Any 2-face of type 4 is commutative because $x_i = x_j = 0 \in V(s \oplus i \oplus j)$. For the face of type 5 we need to show that $x_i = x_j \in V(s \oplus i \oplus j)$. There are three possibilities.

1. $\text{sgn}(v_i) = \text{sgn}(v_j) = -1$. Then $v_i, v_j \in s \oplus i \oplus j$. Without loss of generality we can assume that v_i and v_j are the last vertices in $s \oplus i \oplus j$. The adjacency matrix $A(s \oplus i \oplus j)$ can be represented in the form

$$\begin{pmatrix} A & a^\top & b^\top \\ a & \alpha & \gamma \\ b & \gamma & \beta \end{pmatrix} \quad (3)$$

where $A = A(s)$.


Since $\text{corank } A(s \oplus i) = \text{corank } A(s) - 1$ we have $x_i = 0 \in V(s \oplus i)$. Then the rows of the matrix $A(s \oplus i) = \begin{pmatrix} A & a^\top \\ a & \alpha \end{pmatrix}$ generate the vector $(0 \ 1)$. Hence, the rows of the matrix $A(s \oplus i \oplus j)$ generate the vector $(0 \ 1 \ \delta)$, $\delta \in \mathbb{Z}_2$. If $\delta = 0$ then $x_i = 0 \in V(s \oplus i \oplus j)$ but this is not the case. So $\delta = 1$ and we have the relation $x_i + x_j = 0$ in $V(s \oplus i \oplus j)$.

2. $\text{sgn}(v_i) = -1$, $\text{sgn}(v_j) = 1$. Then $v_i \in s \oplus i \oplus j$ and $v_j \notin s \oplus i \oplus j$. Assume that the adjacency matrix $A(s \oplus i)$ has the form (3) where $A = A(s \oplus j)$.

We have the equality $x_i = 0 \in A(s \oplus i)$. It implies that the vector $(0 \ 1 \ 0)$ is a linear combination of the rows of the adjacency matrix $A(s \oplus i)$. If the coefficient in the linear combination by the row $(b \ \gamma \ \beta)$ is zero then the rows of the matrix $A(s \oplus i \oplus j) = \begin{pmatrix} A & a^\top \\ a & \alpha \end{pmatrix}$ generate the vector $(0 \ 1)$. Thus $x_i = 0 \in V(s \oplus i \oplus j)$ that leads to contradiction. Therefore the coefficient by $(b \ \gamma \ \beta)$ is 1. Then the vector $(b \ \gamma + 1)$ is generated by the rows of the matrix $A(s \oplus i \oplus j)$ that means $x_i + x_j = 0 \in V(s \oplus i \oplus j)$.

3. $\text{sgn}(v_i) = \text{sgn}(v_j) = -1$. Then the matrix $A(s)$ looks like (3) where $A = A(s \oplus i \oplus j)$. The equality $x_i + x_j = 0 \in V(s \oplus i \oplus j)$ corresponds to the relation vector $a + b$. Since $\text{rank } A(s) = \text{rank } A(s \oplus j) = \text{rank} \begin{pmatrix} A & a^\top \\ a & \alpha \end{pmatrix} = \text{rank} \begin{pmatrix} A & a^\top & b^\top \\ a & \alpha & \gamma \end{pmatrix}$ the last row in the matrix $A(s)$ is expressed as a linear combination of the other rows. If the coefficient by the row $(a \ \alpha \ \gamma)$ in the combination is zero then the vector $(b \ \gamma \ \beta)$ depends on the rows of the matrix $\begin{pmatrix} A & a^\top & b^\top \end{pmatrix}$ and the vector b depends on the rows of the matrix A . Hence, $x_j = 0 \in V(s \oplus i \oplus j)$ that is not true. Thus, the coefficients by $(a \ \alpha \ \gamma)$ is not zero so the vector $a + b$ is generated by the rows of the matrix A that means $x_i + x_j = 0 \in V(s \oplus i \oplus j)$.

Any diagram, which has odd edges in both upper-right and left-upper paths, is commutative because the map of any odd edge is zero. After this remark it remains only two nontrivial diagrams (up to symmetry between i and j) with an odd edge:



The diagram of type 6 is commutative because $\dim V(s \oplus i) = \dim V(s \oplus i \oplus j)$ so $x_j = 0 \in V(s \oplus i \oplus j)$.

For the commutativity of the diagram of type 7 we need to prove that $x_i = 0 \in V(s \oplus i \oplus j)$. Assume this is not the case.

Let $\text{sgn}(v_i) = \text{sgn}(v_j) = -1$. Then the adjacency matrix looks like (3). Since $x_j = 0 \in V(s \oplus i)$ we have $\text{rank} \begin{pmatrix} A & a^\top \\ a & \alpha \\ b & \gamma \end{pmatrix} = \text{rank} \begin{pmatrix} A & a^\top \\ a & \alpha \end{pmatrix}$. Then b depends on the rows of $\begin{pmatrix} A \\ a \end{pmatrix}$. But $\text{rank} \begin{pmatrix} A \\ a \end{pmatrix} = \text{rank } A$ hence b depends on the rows of A . On the other hand, we have $x_j \neq 0 \in V(s)$ so $\text{rank} \begin{pmatrix} A \\ b \end{pmatrix} \neq \text{rank } A$. This is contradiction.

The other cases of sign arrangement can be treated in an analogous manner.

This concludes the proof of Proposition 1. $\square$

Definition 2. We call the homology $\overline{\text{Kh}}(G)$ of the complex $(C(G), \partial)$ the *reduced (odd) Khovanov homology* of the labeled simple graph G .

The main theorem of the article states that Khovanov homology is well defined for graph-links.

Theorem 1. *Khovanov homology $\overline{\text{Kh}}(G)$ is invariant under $\Omega_1 - \Omega'_4$ moves.*

Proof. Let G be a labeled graph and $\tilde{G}$ be the graph obtained from G by some Reidemeister move $\Omega_1 - \Omega'_4$.

Invariance under Ω_1 .

Let $\tilde{G}$ be obtained from G by an addition of an isolated labeled vertex v . The complex $C(\tilde{G})$ is isomorphic to the product of complexes $C(G) \otimes C(v)$ where the complex $C(v)$ is equal to

$$\mathbb{Z}_2 \xrightarrow{x \wedge} \bigwedge^* \mathbb{Z}_2 \langle x \rangle$$

if $\text{sgn}(v) = -1$ and

$$\bigwedge^* \mathbb{Z}_2 \langle x \rangle \xrightarrow{x=0} \mathbb{Z}_2$$

if $\text{sgn}(v) = 1$. In any case $H_*(C(v)) = \mathbb{Z}_2 \cdot 1$, where $1 \in H_0(C(v))$ if $\text{sgn}(v) = 1$ and $1 \in H_1(C(v))$ if $\text{sgn}(v) = -1$. Thus, we have

$$\overline{\text{Kh}}(\tilde{G}) = \overline{\text{Kh}}(G) \otimes \overline{\text{Kh}}(v) \cong \overline{\text{Kh}}(G).$$

Invariance under Ω_2 .

Assume that we add the vertices v and w to get the graph $\tilde{G}$ by Ω_2 and $\text{sgn}(v) = 1, \text{sgn}(w) = -1$. Without loss of generality we can write the adjacency matrix $A(\tilde{G})$ in one of this two forms

$$\begin{pmatrix} 0 & 0 & a^\top \\ 0 & 0 & a^\top \\ a & a & A(G) \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} 1 & 1 & a^\top \\ 1 & 1 & a^\top \\ a & a & A(G) \end{pmatrix}.$$

In both cases for every state $s \in \mathcal{V}(G)$ we have the following equalities: $\text{corank } A(G(s)) = \text{corank } A(\tilde{G}(s)), \text{corank } A(\tilde{G}(s \cup \{v\})) = \text{corank } A(\tilde{G}(s \cup \{w\})) = \text{corank } A(\tilde{G}(s \cup \{v, w\}))$

$\{v, w\}) - 1$. These equalities define the type of the upper and left arrows of the complex $C(\tilde{G})$ written in the form

$$\begin{array}{ccc} C_{vw} & \xrightarrow{1} & C_w \\ x_2 \uparrow & & \uparrow \partial \\ C_v & \xrightarrow{\partial} & C. \end{array}$$

Here C_v consists of the chains whose state contains v and does not contain w ; the complexes C, C_w, C_{vw} are defined analogously.

For any state s in C_{vw} let us define a linear function $f : V(s) \rightarrow \mathbb{Z}_2$ by the formula $f(\sum_i \lambda_i x_i) = \lambda_1 + \lambda_2$. The function f is well defined because it equals to zero on any relation: $f(r_i^s) = a_{i1} + a_{i2} = 0$ since $a_{i1} = a_{i2}$. Then $\bigwedge^* V(s) = \bigwedge^* \ker f \oplus x_2 \bigwedge^* V(s)$ and $C_{vw} = X \oplus x_2 C_{vw}$. The subcomplex $X \rightarrow C_w$ is acyclic because the arrow is an isomorphism. Then the homology of $C(\tilde{G})$ coincides with the homology of the quotient complex

$$\begin{array}{ccc} x_2 C_{vw} & & \\ x_2 \uparrow & & \\ C_v & \xrightarrow{\partial} & C. \end{array}$$

The quotient of the obtained complex by the subcomplex C appears to be acyclic too. Thus $C(\tilde{G})$ has the same homology as $C = C(G)$.

Invariance under Ω_3 .

Without loss of generality we can assume that the vertices u, v, w in the third Reidemeister move have the indices 1, 2, 3 in $\mathcal{V}(G) = \mathcal{V}(\tilde{G})$. There are two variants of Ω_3 depending adjacency of v and w and we consider the version where v and w are not adjacent. The proof for the other version is analogous. The adjacency matrices of G and $\tilde{G}$ looks like

$$A(G) = \begin{pmatrix} 0 & 1 & 1 & 0^\top \\ 1 & 0 & 0 & a^\top \\ 1 & 0 & 0 & b^\top \\ 0 & a & b & B \end{pmatrix}, \quad A(\tilde{G}) = \begin{pmatrix} 0 & 0 & 0 & (a+b)^\top \\ 0 & 0 & 0 & a^\top \\ 0 & 0 & 0 & b^\top \\ a+b & a & b & B \end{pmatrix}.$$

Denote $\tilde{V}(s) = V(\tilde{G}(s))$. Then for any $s \in \mathcal{V}(G) \setminus \{u, v, w\}$ we have $V(s) \cong \tilde{V}(s)$, $V(s \oplus v) \cong \tilde{V}(s \oplus v)$, $V(s \oplus w) \cong \tilde{V}(s \oplus w)$, $V(s \oplus v \oplus w) \cong \tilde{V}(s \oplus u \oplus v) \cong \tilde{V}(s \oplus u \oplus w)$, $V(s \oplus u \oplus v \oplus w) \cong \tilde{V}(s \oplus u)$, $V(s \oplus u \oplus v) \cong V(s \oplus u \oplus w) \cong V(s)$ and the correspondent isomorphisms of the exterior algebras are compatible with the differential.

Consider complexes $C(G)$ and $C(\tilde{G})$ in the form of cubes:

$$\begin{array}{ccc}
 & C_{uw} & \longrightarrow C_{uvw} \\
 & \nearrow 1 \uparrow & \nearrow \\
 C_u & \xrightarrow{\quad 1 \quad} C_{uv} & \nearrow \\
 \uparrow x_1 & \uparrow & \uparrow \\
 & C_w & \longrightarrow C_{vw} \\
 & \nearrow & \nearrow \\
 C & \longrightarrow C_v &
 \end{array}
 \qquad
 \begin{array}{ccc}
 & \tilde{C}_{uv} & \longrightarrow \tilde{C}_u \\
 & \nearrow 1 \uparrow & \nearrow \\
 \tilde{C}_{uvw} & \xrightarrow{\quad 1 \quad} \tilde{C}_{uw} & \nearrow \\
 \uparrow x_1 & \uparrow & \uparrow \\
 & \tilde{C}_v & \longrightarrow \tilde{C} \\
 & \nearrow & \nearrow \\
 \tilde{C}_{vw} & \longrightarrow \tilde{C}_w &
 \end{array}$$

For any state s in C_u let us define a linear function $f : V(s) \rightarrow \mathbb{Z}_2$ by the formula $f(\sum_i \lambda_i x_i) = \lambda_1$. The function is well defined and there are decompositions $\bigwedge^* V(s) = \bigwedge^* \ker f \oplus x_1 \bigwedge^* V(s)$ and $C_u = X \oplus x_1 C_u$. Consider the following subcomplex

$$\begin{array}{ccc}
 & C_{uw} & \longrightarrow C_{uvw} \\
 & \nearrow 1 \uparrow & \nearrow \\
 X & \xrightarrow{\quad 1 \quad} C_{uv} & \nearrow \\
 & \uparrow & \uparrow \\
 & C_w & \longrightarrow C_{vw} \\
 & \nearrow & \nearrow \\
 & C_v &
 \end{array}$$

The quotient complex $C \rightarrow x_1 C_u$ is acyclic so the homology of $C(G)$ is isomorphic to the homology of the subcomplex. This subcomplex contains the acyclic subcomplex $X \rightarrow \partial(X)$. The maps $X \rightarrow C_{uv}$ and $X \rightarrow C_{uw}$ are isomorphisms, so after factorization we obtain the complex

$$\begin{array}{ccc}
 C_{uw} & \longrightarrow & C_{uvw} \\
 \uparrow & \searrow & \nearrow \\
 & C_{uv} & \\
 \uparrow & \nearrow & \uparrow \\
 C_w & \longrightarrow & C_{vw} \\
 & \nearrow & \nearrow \\
 & C_v &
 \end{array}$$

where the spaces C_v and C_w are identified.

The reasonings analogous to the reasonings above reduce $C(\tilde{G})$ to the complex (for a state s in $\tilde{C}_{uvw}$ we should define the function $f : \tilde{V}(s) \rightarrow \mathbb{Z}_2$ by the formula $f(\sum_i \lambda_i x_i) = \lambda_1 + \lambda_2 + \lambda_3$).

$$\begin{array}{ccc}
 \tilde{C}_{uv} & \longrightarrow & \tilde{C}_u \\
 \uparrow & \searrow & \nearrow \\
 & \tilde{C}_{uw} & \\
 \uparrow & \nearrow & \uparrow \\
 \tilde{C}_v & \longrightarrow & \tilde{C} \\
 & \nearrow & \nearrow \\
 & \tilde{C}_w &
 \end{array}$$

Both complexes are isomorphic to the complex $C_v \oplus C_w \rightarrow C \oplus C_{vw} \rightarrow C_{uvw}$. Thus $C(G)$ and $C(\tilde{G})$ have the same homology.

Invariance under Ω_4 .

Let the vertices u and v of the move Ω_4 have the indices p and q in $V(G) = V(\tilde{G})$. The coefficients of adjacency matrices of $A(G) = (a_{ij})$ and $A(\tilde{G}) = (\tilde{a}_{ij})$ are connected by the formula

$$\tilde{a}_{ij} = \begin{cases} a_{ij} + a_{ip}a_{jq} + a_{iq}a_{jp}, & \{i, j\} \cap \{p, q\} = \emptyset, \\ a_{ij}, & \{i, j\} \cap \{p, q\} \neq \emptyset. \end{cases}$$

Consider the map ϕ acting on the states by the formula

$$\phi(s) = \begin{cases} s \cup \{u, v\}, & \{u, v\} \cap s = \emptyset, \\ s \setminus \{u, v\}, & \{u, v\} \cap s = \{u, v\}, \\ s, & \{u, v\} \cap s \neq \emptyset, \{u, v\} \end{cases}$$

together with the linear maps $\Phi : V(s) \rightarrow \tilde{V}(\phi(s))$ defined by the formula

$$\Phi(x_i) = \begin{cases} x_i, & i \neq p, q, \\ x_q, & i = p, \\ x_p, & i = q. \end{cases}$$

One can check that the map Φ defines an isomorphism of the vector spaces and after the natural extension to an isomorphism of external algebras $\bigwedge^* V(s) \rightarrow \bigwedge^* \tilde{V}(\phi(s))$ for each state s it determines an isomorphism of the graded linear spaces $\Phi : C(G) \rightarrow C(\tilde{G})$. Φ is a chain map. Thus the complexes $C(G)$ and $C(\tilde{G})$ are isomorphic as well as their homology.

Invariance under Ω'_4 .

Let the vertex v of the move Ω'_4 have the index p . The coefficients of adjacency matrices of $A(G) = (a_{ij})$ and $A(\tilde{G}) = (\tilde{a}_{ij})$ are connected by the formula

$$\tilde{a}_{ij} = \begin{cases} a_{ij} + a_{ip}a_{jp}, & i, j \neq p, \\ a_{ip}, & j = p, \\ a_{pj}, & i = p, \end{cases}$$

Consider the map $\phi : C(G) \rightarrow C(\tilde{G})$ which acts on the states by the formula

$$\phi(s) = s \oplus p$$

and the linear maps $\Phi : V(s) \rightarrow \tilde{V}(\phi(s))$ defined on the generators by the formula $\Phi(x_i) = x_i$, $1 \leq i \leq n$. Then Φ induces a well defined isomorphism of the complexes $C(G)$ and $C(\tilde{G})$ so the homologies of the complexes coincide. $\square$

Corollary 2. *Khovanov homology $\overline{\text{Kh}}(G)$ is an invariant of graph-links.* $\square$

4 Jones polynomial of graph-knots and Khovanov homology

Let G be a simple labeled graph. The complex $C(G)$ has two grading: the homological grading M_0 and the algebraic grading $\deg$ of graded algebras $\bigwedge^* V(s)$. The differential is not homogeneous with respect to $\deg$ but it is compatible with the grading Q_0 where for element any $u \in \bigwedge^r(s)$ we define $Q_0(u) = \dim_{\mathbb{Z}_2} V(s) - 2r + M_0(s)$. The differential increases the grading M_0 and leaves the grading Q_0 unchanged. Then

$$\overline{\text{Kh}}(G) = \bigoplus_{m,q \in \mathbb{Z}} \overline{\text{Kh}}(G)_{(m,q)}.$$

The following definition is due to Ilyutko and Manturov [4].

Definition 3. Let G be a simple labeled graph with n vertices. The *Kauffman bracket polynomial* of G is defined as

$$\langle G \rangle(a) = \sum_{s \subset \mathcal{V}(G)} a^{\alpha(s) - \beta(s)} (-a^2 - a^{-2})^{\text{corank } A(s)},$$

where $\alpha(s) = \#\{v \in s \mid \text{sgn}(v) = -1\} + \#\{v \notin s \mid \text{sgn}(v) = 1\}$ and $\beta(s) = n - \alpha(s)$.

It appears that the Poincaré polynomial of the bigraded Khovanov homology of the graph G coincides with its Kauffman bracket up to shift of the gradings.

Theorem 2.

$$\sum_{m,q \in \mathbb{Z}} (-1)^m \dim_{\mathbb{Z}_2} \overline{\text{Kh}}(G)_{(m,q)} \cdot t^q = (it^{1/2})^{-n} \langle G \rangle(it^{1/2}).$$

Proof. The left term of the equality coincides with the Poincaré polynomial of the chain complex $C(G)$. Each state $s \subset \mathcal{V}(G)$ has homological grading $\alpha(s)$ and the corresponding chain space $\bigwedge^* V(s)$ contributes $(-1)^{\alpha(s)} (t + t^{-1})^{\dim V(s)} t^{\alpha(s)}$ to the Poincaré polynomial. Since $\dim V(s) = \text{corank } A(s)$ and $\alpha(s) = \frac{1}{2}(\alpha(s) - \beta(s) - n)$ the sum over all states yields the right term of the equality. $\square$

The gradings M_0, Q_0 of the Khovanov complex are not preserved by the Reidemeister moves as shown in the following table.

	M_0	Q_0
Ω_1^+	0	-1
Ω_1^-	1	1
Ω_2	1	0
Ω_3	0	0
Ω_4	0	0
Ω'_4	0	0

(4)

Here $\Omega_1^\pm$ denotes the adding an isolated vertex with the label ± 1 and Ω_2 denotes the adding two vertices. The entries of the cells are the shifts of the gradings after the corresponding move.

Thus the groups $\overline{\text{Kh}}(G)_{(m,q)}$ are not invariants of graph-links. Nevertheless we can normalize the gradings for the graph-knots.

Definition 4 ([4]). A graph-link G is called a *graph-knots* if $\text{corank}(A(G) + E) = 0$ where E is the identity matrix.

The condition $\text{corank}(A(G) + E) = 0$ survives after Reidemeister moves so the definition is correct.

Definition 5 ([4]). *Writhe number* of a graph G with n vertices is the sum

$$w(G) = \sum_{i=1}^n (-1)^{\text{corank } B_i(G)} \text{sgn } v_i,$$

where $B_i(G) = A(G) + E + E_{ii}$ and the only non-zero element of E_{ii} lies in the i -th row and i -th column.

Writhe number is invariant under the moves $\Omega_2 - \Omega_4$. The move $\Omega_1^\pm$ changes the writhe number by ∓ 1 . Using this fact and the table (4) we construct two gradings which are invariant under the Reidemeister moves

$$M = M_0 - \frac{n(G) + w(G)}{2}, \quad Q = Q_0 + \frac{n(G) + 3w(G)}{2}$$

where $n(G)$ is the number of vertices in G .

Let us denote $\overline{\text{Kh}}_{m,q}(G)$ to be the homogeneous part of $\overline{\text{Kh}}(G)$ with the gradings $M = m$ and $Q = q$. The groups $\overline{\text{Kh}}_{m,q}(G)$ are invariants of graph-knots.

Writhe number allows to introduce another invariant of graph-knots.

Definition 6 ([4]). *Jones polynomial* of a graph-knot G is defined as

$$X(G)(a) = (-a)^{-3w(G)} \langle G \rangle(a).$$

Theorem 2 leads to the following statement.

Theorem 3. *The bigraded Khovanov homology of graph-knots $\overline{\text{Kh}}_{m,q}(G)$ categorifies the Jones polynomial in the sense that*

$$\sum_{m,q \in \mathbb{Z}} (-1)^m \dim_{\mathbb{Z}_2} \overline{\text{Kh}}_{m,q}(G) \cdot t^q = X(G)(it^{1/2}).$$

The author is grateful to V. Manturov and D. Ilyutko for useful discussions.

References

- [1] J. Bloom, Odd khovanov homology is mutation invariant // Math. Res. Lett. **17**(1) (2010), 1–10.
- [2] S. Chmutov, S. Lando, Mutant knots and intersection graphs // Alg. Geom. Topology **7** (2007), 1579–1598.
- [3] D. Ilyutko, Framed 4-Valent Graphs: Euler Tours, Gauss Circuits and Rotating Circuits // arXiv:Math.CO/0911.5504.
- [4] D. Ilyutko, V. Manturov, Introduction to graph-link theory // Journal of Knot Theory and Its Ramifications, **18**(6) (2009), 791–823.
- [5] D. Ilyutko, V. Manturov, Graph links // Dokl. Akad. Nauk 428 (2009), no. 5, 591–594 (Russian).
- [6] P. Ozsváth, J. Rasmussen, Z. Szabó, Odd Khovanov homology // www.arxiv.org/math-qa/0710.4300.
- [7] L. Traldi, L. Zulli, A bracket polynomial for graphs // arXiv:math.GT/0808.3392.
- [8] S. Wehrli, Khovanov homology and Conway mutation // arXiv:math.GT/0301.312.